

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody Goat Polyclonal, Biotin

Catalog # ASR2121

Specification

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Product Information

Description

- Host Conjugate Target Species Clonality Application Application Note
- Physical State Host Isotype Target Isotype Buffer

Immunogen Species of Origin Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Conjugated Goat Biotin Ferret Polyclonal ,1,10,15, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Lyophilized laG laA 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Ferret IgA alpha heavy chain **Ferret** 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

Anti-FERRET IqA(GOAT) Antibody Biotin

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

Anti-FERRET IgA (alpha chain) (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Ferret IgA coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Ferret IgA and Ferret Serum. No reaction was observed against Ferret IgM or Ferret IgG.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Images

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Background

Ferret IgA whole molecule antibody generated in goat detects specifically ferret IgA whole molecule. Anti-Human IgA is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays