10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody
**Goat Polyclonal, Rhodamine (TRITC)**
**Catalog # ASR2253**

## Specification

### Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody - Product Information

| | |
|---|---|
| Description | **Anti-MONKEY IgA (alpha chain) (GOAT) Antibody Rhodamine Conjugated** |
| Host | **Goat** |
| Conjugate | **Rhodamine (TRITC)** |
| FP Value | **2.9 moles Rhodamine (TRITC) per mole of IgG** |
| Target Species | **Monkey** |
| Clonality | **Polyclonal** |
| Application | **,3,4,** |
| Application Note | **FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500** |
| Physical State | **Lyophilized** |
| Host Isotype | **IgG** |
| Target Isotype | **IgA** |
| Buffer | **0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2** |
| Immunogen | **Monkey IgA alpha heavy chain** |
| Reconstitution Volume | **1.0 mL** |
| Reconstitution Buffer | **Restore with deionized water (or equivalent)** |
| Stabilizer | **10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free** |
| Preservative | **0.01% (w/v) Sodium Azide** |

### Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody - Additional Information

**Shipping Condition**
Ambient

**Purity**
This product was prepared from monospecific antiserum by immunoaffinity chromatography using Monkey IgA coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Monkey IgA and Monkey Serum. No reaction was observed against Monkey IgG or Monkey IgM. Specificity was confirmed by ELISA at less than 1% cross reactivity against other Monkey heavy or light chain isotypes.

**Storage Condition**
Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid.  Dilute only prior to immediate use.

**Precautions Note**
This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody - Protein Information

## Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody - Images

| Product | Gt-a-Monkey IgG | Gt-a-Monkey IgA | Gt-a-Monkey IgM |
|---|---|---|---|
| Rhesus Serum | + | + | + |
| Baboon Serum | + | + | + |
| Cynomologus Serum | + | + | + |
| Rhesus IgG | + | | |
| Human IgA | | + | |
| Human IgM | | | + |

This table displays additional reactivity among various species of serum and immunoglobulin. A (+) indicates antibody reactivity to the corresponding target.

## Anti-Monkey IgA (alpha chain) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.