

MOUSE IgG1 Isotype control

Monoclonal MG1K IgG1 , Unconjugated Catalog # ASR2265

Specification

MOUSE IgG1 Isotype control - Product Information

Description Conjugate Clonality Application Application Note Physical State Host Isotype Buffer

Species of Origin Stabilizer Preservative MOUSE IgG1 Kappa (κ) isotype control Unconjugated Monoclonal ,4, FlowCytometry 1:1000-1:5000 Liquid (sterile filtered) IgG1 0.02 M Potassium Phosphate, 0.5 M Sodium Chloride, pH 7.2 Mouse None 0.01% (w/v) Sodium Azide

MOUSE IgG1 Isotype control - Additional Information

Shipping Condition Wet Ice

Purity

Mouse Isotype control has been prepared from concentrated cell culture supernatant by immunoaffinity chromatography using protein A. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to mouse IgG2a, IgG2b, IgG3, IgM, and IgA. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse IgG and anti-Mouse serum. Light and heavy chain composition has been confirmed..

Storage Condition

Store vial at 4° C prior to opening. This product is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

MOUSE IgG1 Isotype control - Protein Information

MOUSE IgG1 Isotype control - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Western Blot</u>



- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

MOUSE IgG1 Isotype control - Images

MOUSE IgG1 Isotype control - Background

Mouse isotype controls are used in flow cytometry, western blot and ELISA and differentiate between immunoglobulin classes and subclasses. Isotype controls allow for the genetic variations or differences in the constant regions of the heavy and light chains. In mouse there are six relevant heavy chain isotypes and two light chain isotypes: heavy chain a - IgA, ? - IgG 1, 2a, 2b, 3 and μ - IgM, light chain ? and ?.