

# Mouse IgG3 Kappa (κ) isotype Control

Monoclonal MG3K IgG3 , Unconjugated Catalog # ASR2271

#### **Specification**

## Mouse IgG3 Kappa (κ) isotype Control - Product Information

Description MOUSE IgG3 Kappa (κ) isotype control

Conjugate Unconjugated Clonality Monoclonal

Application ,4,

Application Note FlowCytometry 1:1000-1:5000
Physical State Liquid (sterile filtered)

Physical State
Host Isotype
Liquid (sterile fi

Buffer 0.02 M Potassium Phosphate, 0.5 M

Sodium Chloride, pH 7.2

Species of Origin
Stabilizer

Mouse
None

Preservative 0.01% (w/v) Sodium Azide

## Mouse IgG3 Kappa (κ) isotype Control - Additional Information

# **Shipping Condition**

Wet Ice

#### **Purity**

Mouse Isotype control has been prepared from concentrated cell culture supernatant by immunoaffinity chromatography using protein A. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse IgG and anti-Mouse serum. Light and heavy chain composition confirmed by ELISA.

## **Storage Condition**

Store vial at 4° C prior to opening. This product is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Mouse IgG3 Kappa (κ) isotype Control - Protein Information

#### Mouse IgG3 Kappa (κ) isotype Control - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides





• Dot Blot

- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# Mouse IgG3 Kappa (κ) isotype Control - Images

# Mouse IgG3 Kappa (κ) isotype Control - Background

Mouse isotype controls are used in flow cytometry, western blot and ELISA and differentiate between immunoglobulin classes and subclasses. Isotype controls allow for the genetic variations or differences in the constant regions of the heavy and light chains. In mouse there are six relevant heavy chain isotypes and two light chain isotypes: heavy chain a - IgA, ? - IgG 1, 2a, 2b, 3 and  $\mu$  - IgM, light chain ? and ?.