

Anti-Guinea Pig IgG F(c) (Biotin Conjugated) Secondary Antibody Rabbit Polyclonal, Biotin Catalog # ASR2331

Specification

Anti-Guinea Pig IgG F(c) (Biotin Conjugated) Secondary Antibody - Product Information

Description

- Host Conjugate Target Species Clonality Application Application Note
- Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-GUINEA PIG IgG F(c) (RABBIT) **Antibody Biotin Conjugated Rabbit** Biotin **Guinea** Pig Polyclonal ,1,2,10, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000 Lyophilized laG laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Guinea Pig IgG F(c) fragment 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

Anti-Guinea Pig IgG F(c) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Guinea Pig IgG, Guinea Pig IgG F(c) and Guinea Pig Serum. No reaction was observed against Guinea Pig IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Guinea Pig IgG F(c) (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Guinea Pig IgG F(c) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Guinea Pig IgG F(c) (Biotin Conjugated) Secondary Antibody - Images