

Anti-Horse IgG F(ab')₂ (Biotin Conjugated) Secondary Antibody
Goat Polyclonal, Biotin
Catalog # ASR2336**Specification****Anti-Horse IgG F(ab')₂ (Biotin Conjugated) Secondary Antibody - Product Information**

Description	Anti-HORSE IgG F(ab')₂ (GOAT) Antibody Biotin Conjugated
Host	Goat
Conjugate	Biotin
Target Species	Horse
Clonality	Polyclonal
Application	,1,2,10, ELISA 1:20,000-1:100,000;Western Blot
Application Note	1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG F(ab')₂
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Horse IgG F(ab')₂ fragment
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-Horse IgG F(ab')₂ (Biotin Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Horse IgG, Horse IgG F(ab')₂ and Horse Serum. No reaction was observed against Horse IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Horse IgG F(ab')₂ (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Horse IgG F(ab')₂ (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Horse IgG F(ab')₂ (Biotin Conjugated) Secondary Antibody - Images