

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody Rabbit Polyclonal, Rhodamine (TRITC)

Catalog # ASR2368

Specification

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Product Information

Description

Host Conjugate FP Value

Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-SWINE IgG F(c) (RABBIT) Antibody **Rhodamine Conjugated** Rabbit **Rhodamine (TRITC)** 2.0 moles Rhodamine (TRITC) per mole of laG Swine Polyclonal ,3,4, FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2.500 Lyophilized lgG lgG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Swine IgG F(c) fragment 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Swine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Swine IgG, Swine IgG F(c) and Swine Serum. No reaction was observed against Swine IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Swine IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.