

**F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody**  
**Donkey Polyclonal, Biotin**  
**Catalog # ASR2383****Specification****F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Product Information**

Description	<b>F(ab')<sub>2</sub> Anti-GOAT IgG [H&amp;L] (DONKEY) Antibody Biotin Conjugated Min X Ch GP Ham Hs Hu Ms Rb &amp; Rt Serum Proteins</b>
Host	<b>Donkey</b>
Conjugate	<b>Biotin</b>
Target Species	<b>Goat</b>
Clonality	<b>Polyclonal</b>
Application	<b>,1,10,15,</b>
Application Note	<b>ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:5,000;Immunocytochemistry 1:500-1:3,000</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG F(ab')<sub>2</sub></b>
Target Isotype	<b>IgG (H&amp;L)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Goat IgG whole molecule</b>
Reconstitution Volume	<b>500 µL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information****Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Pepsin or anti-Donkey IgG F(c), Chicken, Guinea Pig, Syrian Hamster, Horse, Human, Mouse, Rabbit, or Rat Serum Proteins.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information****F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Images****F(ab')<sub>2</sub> Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Background**

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.