

F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody
Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2388**Specification****F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody**
- Product Information

Description	F(ab')₂ Anti-HUMAN IgG F(c) (GOAT) Antibody Peroxidase Conjugated Min X Bv Hs Ms & Rt Serum Proteins
Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Human
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:130,000;Western Blot 1:1,000-1:5,000;Immunochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG F(ab')₂
Target Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Human IgG F(c) fragment
Reconstitution Volume	500 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody
- Additional Information**Shipping Condition**

Ambient

Purity

F(ab')₂ Anti-Human IgG Fc was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Human IgG F(ab')₂ or Bovine, Horse, Mouse and Rat Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Images**F(ab')₂ Anti-Human IgG F(c) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Background**

F(ab')₂ Anti-Human F(c) antibody was generated in goat and detects specifically Human IgG F(c) fragment. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.