

F(ab')2 Anti-Human IgM Fc5µ (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish) Catalog # ASR2390

Specification

F(ab')2 Anti-Human IgM Fc5 μ (Peroxidase Conjugated) Secondary Antibody - Product Information

Description	F(ab')2 Anti-HUMAN IgM Fc5 μ (GOAT) Antibody Peroxidase Conjugated
Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Human
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:20,000;Western Blot
	1:2,000-1:10,000;Immunochemistry
	1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	lgG F(ab')2
Target Isotype	lgM Fc5 μ
Buffer	0.02 M Potassium Phosphate, 0.15 M
	Sodium Chloride, pH 7.2
Immunogen	Human IgM Fc5 μ fragment
Reconstitution Volume	500 μL
Reconstitution Buffer	Restore with deionized water (or
	equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) -
	Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate

F(ab')2 Anti-Human IgM Fc5µ (Peroxidase Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgM and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Human IgG or Human IgA. Specificity was confirmed by ELISA at less than 1% cross reactivity against other human heavy or light chain isotypes.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Human IgM Fc5 μ (Peroxidase Conjugated) Secondary Antibody - Protein Information

F(ab')2 Anti-Human IgM Fc5µ (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

F(ab')2 Anti-Human IgM Fc5µ (Peroxidase Conjugated) Secondary Antibody - Images