10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Goat IgG F(ab')2 (Agarose Conjugated)

**Catalog # ASR2415**

## Specification

### Goat IgG F(ab')2 (Agarose Conjugated) - Product Information

| | |
|---|---|
| Description | **GOAT IgG F(ab')2 fragment Agarose Conjugated** |
| Conjugate | **Unconjugated** |
| Physical State | **Suspension of agarose beads** |
| Buffer | **0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2** |
| Species of Origin | **Goat** |
| Stabilizer | **None** |
| Preservative | **0.01% (w/v) Sodium Azide** |

### Goat IgG F(ab')2 (Agarose Conjugated) - Additional Information

**Shipping Condition**
Wet Ice

**Purity**
This product is an purified Goat IgG isolated from normal Goat Serum digested by the enzyme pepsin and purified by chromatography coupled to activated agarose.  Sufficient antibody capacity is provided to bind a minimum of 5.0 mg of anti-Goat IgG.

**Storage Condition**
Store vial at 4° C prior to opening. DO NOT FREEZE.

**Precautions Note**
This product is for research use only and is not intended for therapeutic or diagnostic applications.

### Goat IgG F(ab')2 (Agarose Conjugated) - Protein Information


### Goat IgG F(ab')2 (Agarose Conjugated) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Goat IgG F(ab')2 (Agarose Conjugated) - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




SDS-PAGE of Goat IgG F(ab')2 Fragment Agarose Conjugated . Lane M: 5 µL Opal Prestained Marker . Lane 1: Reduced Goat IgG Whole Molecule . Lane 2: Reduced Goat IgG F(c) Fragment . Lane 3: Reduced Goat IgG F(ab')2 Fragment Agarose Conjugated . Lane 4: Reduced Goat IgM Whole Molecule . Load: 1 µg for IgG, F(c) and F(ab')2; 3 µg for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.