

## RAT IgG2b Kappa (κ) isotype control

Monoclonal RG2bK IgG2b kappa , Unconjugated Catalog # ASR2538

### **Specification**

## RAT IgG2b Kappa (κ) isotype control - Product Information

Description RAT IgG2b Kappa (κ) isotype control

Conjugate Unconjugated Clonality Monoclonal

Application ,4,

Application Note FlowCytometry 1:1000-1:5000
Physical State Liquid (sterile filtered)

Host Isotype IgG2b

Buffer 0.02 M Potassium Phosphate, 0.5 M

Sodium Chloride, pH 7.2

Species of Origin
Stabilizer
Rat
None

Preservative 0.01% (w/v) Sodium Azide

### RAT IgG2b Kappa (κ) isotype control - Additional Information

## **Shipping Condition**

Wet Ice

#### **Purity**

RAT IgG2b Kappa isotype control has been prepared from concentrated cell culture supernatant by immunoaffinity chromatography using protein G. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to rat IgG1a, IgG1b, IgG3, IgM, and IgA. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rat IgG and anti-Rat serum. Light and heavy chain composition has been confirmed.

## **Storage Condition**

Store vial at 4° C prior to opening. This product is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

# **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

# RAT IgG2b Kappa (κ) isotype control - Protein Information

### RAT IgG2b Kappa (κ) isotype control - Protocols

Provided below are standard protocols that you may find useful for product applications.

Western Blot



Tel: 858.875.1900 Fax: 858.875.1999



- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

RAT IgG2b Kappa (κ) isotype control - Images

RAT IgG2b Kappa (κ) isotype control - Background

RAT IgG2b Kappa isotype control is used in flow cytometry, western blot and ELISA and differentiate between immunoglobulin classes and subclasses. Isotype controls allow for the genetic variations or differences in the constant regions of the heavy and light chains. In Rat there are six relevant heavy chain isotypes and two light chain isotypes: heavy chain a - IgA, ? - IgG 1, 2a, 2b, 2c and  $\mu$  - IgM, light chain ? and ?.