10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Goat IgM
Catalog # ASR2561

Specification

Goat IgM - Product Information

Description	**GOAT IgM whole molecule**
Conjugate	**Unconjugated**
Physical State	**Liquid (sterile filtered)**
Host Isotype	**IgM**
Buffer	**0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0**
Species of Origin	**Goat**
Stabilizer	**10% (v/v) Glycerol**
Preservative	**0.1% (w/v) Sodium Azide**

Goat IgM - Additional Information

Shipping Condition
Wet Ice

Purity
This product was prepared from normal serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum and anti-Goat IgM (μ chain specific). No reaction was observed against anti-Goat IgG F(c). Some light chain cross reactivity will occur with anti-Goat IgG.

Storage Condition
Store vial at 4° C prior to opening. This product is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

Goat IgM - Protein Information

Goat IgM - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat IgM - Images



SDS-PAGE of Goat IgM Whole Molecule . Lane M: 5 µL Opal Prestained Marker . Lane 1: Reduced Goat IgG Whole Molecule . Lane 2: Reduced Goat IgG F(c) Fragment . Lane 3: Reduced Goat IgG F(ab) Fragment . Lane 4: Reduced Goat IgM Whole Molecule . Load: 1 µg for IgG, F(c) and F(ab); 3 µg for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.