

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody

Rabbit Polyclonal, Texas Red® Catalog # ASR2618

Specification

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Product Information

Description Anti-BOVINE IgG F(c) (RABBIT) Antibody

Texas Red™ Conjugated

Host Rabbit Conjugate Texas Red®

FP Value 3.5 moles Texas Red® per mole of IgG

Target Species
Clonality
Application
Bovine
Polyclonal
,3,4,

Application Note FLISA 1:10,000-1:50,000; IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2,500 Lyophilized

Physical State
Host Isotype
Target Isotype
IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2
Bovine IgG F(c) fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/ml Polyethylene Glycol (PEG-8000)

Preservative 0.01% (w/v) Sodium Azide

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Immunogen

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG, Bovine IgG F(c) and Bovine Serum. No reaction was observed against Bovine IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Protein Information

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Images

Anti-Bovine IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.