

F(ab')2 Fragment of Affinity Purified

Anti-GOLDEN SYRIAN & ARMENIAN

# F(ab')2 Anti-Fragment Of Hamster IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish) Catalog # ASR2821

Specification

## F(ab')2 Anti-Fragment Of Hamster IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description

	HAMSTER IGG (H&L) (GOAT) Antibody
	Peroxidase Conjugated (Min X MOUSE and
	RAT Serum Proteins)
Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Armenian and Golden Syrian Hamster
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:20,000;Western Blot
	1:2,000-1:10,000;Immunochemistry
	1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	lgG F(ab')2
Target Isotype	lgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M
	Sodium Chloride, pH 7.2
Immunogen	Golden Syrian and Armenian Hamster IgG
	whole molecules
Reconstitution Volume	500 μL
Reconstitution Buffer	Restore with deionized water (or
	equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) -
	Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT
	add Sodium Azide!

## F(ab')2 Anti-Fragment Of Hamster IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian and Armenian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Golden Syrian and Armenian Hamster IgG and Golden Syrian and Armenian Hamster Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) and Mouse or Rat Serum Proteins.



#### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note** This product is for research use only and is not intended for therapeutic or diagnostic applications.

### F(ab')2 Anti-Fragment Of Hamster IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information

## F(ab')2 Anti-Fragment Of Hamster IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

F(ab')2 Anti-Fragment Of Hamster IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Images