

Swine IgG F(c) Biotin

Catalog # ASR2837

Specification

Swine IgG F(c) Biotin - Product Information

Description SWINE IgG F(c) fragment Biotin

Conjugated
Conjugate
Physical State
Host Isotype

Conjugated
Biotin
Lyophilized
Lyophilized
IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Species of Origin
Reconstitution Volume

Swine
1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Swine IgG F(c) Biotin - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from normal serum delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Swine IgG, anti-Swine IgG F(c) and anti-Swine Serum. No reaction was observed against anti-Swine IgG F(ab')2 or anti-Papain.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Swine IgG F(c) Biotin - Protein Information

Swine IgG F(c) Biotin - Protocols

Provided below are standard protocols that you may find useful for product applications.



Tel: 858.875.1900 Fax: 858.875.1999



- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Swine IgG F(c) Biotin - Images