

Guinea Pig IgM
Catalog # ASR2897**Specification**

Guinea Pig IgM - Product Information

Description	GUINEA PIG IgM whole molecule
Conjugate	Unconjugated
Physical State	Liquid (sterile filtered)
Host Isotype	IgM
Buffer	0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0
Species of Origin	Guinea Pig
Preservative	0.1% (w/v) Sodium Azide

Guinea Pig IgM - Additional Information**Shipping Condition**

Wet Ice

Purity

Guinea Pig IgM whole molecule was prepared from normal serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Guinea Pig IgM whole molecule was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Guinea Pig Serum and anti-Guinea Pig IgM (μ chain specific). No reaction was observed against anti-Guinea Pig IgG F(c). Some light chain cross reactivity will occur with anti-Guinea Pig IgG.

Storage Condition

Store vial at 4° C prior to opening. Guinea Pig IgM whole molecule is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


Guinea Pig IgM - Protein Information**Guinea Pig IgM - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Guinea Pig IgM - Images



SDS-Page of Guinea Pig IgM Whole Molecule. Lane 1: Guinea Pig IgM, Non-Reduced. Lane 2: Guinea Pig IgM, Reduced. Load: 1.0 μ g per lane. Predicted/Observed size-Predicted/Observed size - Non-Reduced: 900 kDa (Pentamer), 900 kDa (Molecule larger than can pass through gel), Reduced: 78 and 25 kDa, 75 and 25 kDa.

Guinea Pig IgM - Background

Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum.