

**Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody**  
**Goat Polyclonal, Peroxidase (Horseradish)**  
**Catalog # ASR3006****Specification****Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody - Product Information**

|                       |   |
|-----------------------|---|
| Description           | <b>Anti-MONKEY IgA (alpha chain) (GOAT) Antibody Peroxidase Conjugated</b>  |
| Host                  | <b>Goat</b>   |
| Conjugate             | <b>Peroxidase (Horseradish)</b>   |
| Target Species        | <b>Monkey</b>   |
| Clonality             | <b>Polyclonal</b>   |
| Application           | <b>,1,10,15,</b>  |
| Application Note      | <b>ELISA 1:10,000-1:150,000;Western Blot 1:1,000-1:10,000;Immunochemistry 1:500-1:2,500</b>                           |
| Physical State        | <b>Lyophilized</b>  |
| Host Isotype          | <b>IgG</b>  |
| Target Isotype        | <b>IgA</b>  |
| Buffer                | <b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>   |
| Immunogen             | <b>Anti-Monkey IgA (alpha chain) was produced by repeated immunization with monkey IgA alpha heavy chain in goat.</b> |
| Reconstitution Volume | <b>1.0 mL</b>   |
| Reconstitution Buffer | <b>Restore with deionized water (or equivalent)</b>   |
| Stabilizer            | <b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>   |
| Preservative          | <b>0.01% (w/v) Thimerosal</b>   |

**Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody - Additional Information****Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Monkey IgA coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Monkey IgA and Monkey Serum. No reaction was observed against Monkey IgM or Monkey IgG.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

## Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


## Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody - Protein Information

## Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody - Images



Western Blot of Peroxidase conjugated Goat anti-MONKEY IgA alpha antibody. Lane 1: Human IgA. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Peroxidase goat secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 60 kDa for Monkey IgA. Other band(s): none.

| Product          | Gt-a-Monkey IgG | Gt-a-Monkey IgA | Gt-a-Monkey IgM |
|------------------|-----------------|-----------------|-----------------|
| Rhesus Serum     | +               | +               | +               |
| Baboon Serum     | +               | +               | +               |
| Cynomolgus Serum | +               | +               | +               |
| Rhesus IgG       | +               |                 |                 |
| Human IgA        |                 | +               |                 |
| Human IgM        |                 |                 | +               |

This table displays additional reactivity among various species of serum and immunoglobulin. A (+) indicates antibody reactivity to the corresponding target.

**Anti-MONKEY IgA alpha (Peroxidase Conjugated) Secondary Antibody - Background**

Anti-Monkey IgA (alpha chain) peroxidase conjugated antibody generated in goat detects specifically Monkey IgA. This secondary peroxidase conjugated antibody anti-Monkey is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.