10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody

Rabbit Polyclonal, Unconjugated
Catalog # ASR3490

Specification

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody - Product Information

Description	F(ab')2 Anti-SHEEP IgG (H&L) (RABBIT) Antibody
Host	Rabbit
Conjugate	Unconjugated
Target Species	Sheep
Clonality	Polyclonal
Application	,1,2,10,
Application Note	ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG F(ab')2
Target Isotype	IgG (H&L)
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Sheep IgG whole molecule
Reconstitution Volume	2.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	None
Preservative	None

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition
Ambient

Purity
This product is a F(ab')2 fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Sheep IgG and Sheep Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.

Storage Condition
Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody - Protein Information

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody - Images

F(ab')2 Anti-Sheep IgG (H&L) Secondary Antibody - Background

F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.