

Anti-Human IgG F(c) Secondary Antibody
Goat Polyclonal, Unconjugated
Catalog # ASR3528**Specification**

Anti-Human IgG F(c) Secondary Antibody - Product Information

Description	Anti-HUMAN IgG F(c) (GOAT) Antibody
Host	Goat
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	,1,2,10,
Application Note	ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG F(c)
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Human IgG F(c) fragment
Reconstitution Volume	5.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Preservative	None

Anti-Human IgG F(c) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

Anti-HUMAN IgG F(c) (GOAT) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab')₂.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgG F(c) Secondary Antibody - Protein Information

Anti-Human IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human IgG F(c) Secondary Antibody - Images

Anti-Human IgG F(c) Secondary Antibody - Background

Anti-Human IgG F(ab')₂ fragment antibody generated in goat detects specifically Human IgG F(ab')₂ fragment. Anti-Human IgG F(ab')₂ antibody is ideal for investigators Involved in Serum Protein Component Protein research.