

Anti-Rabbit IgG (H&L) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR3537

Specification

Anti-Rabbit IgG (H&L) Secondary Antibody - Product Information

Description Host Conjugate Target Species Reactivity Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen

Reconstitution Volume Reconstitution Buffer

Stabilizer Preservative Anti-RABBIT IgG (H&L) (GOAT) Antibody Goat Unconjugated Rabbit Rabbit Polyclonal ,1,10,15, ELISA 1:50,000;Western Blot 1:5,000-1:25,000;Immunochemistry 1:1,000-1:5,000 Lyophilized laG IgG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Anti-Rabbit IgG whole molecule was produced by repeated immunization with Rabbit IgG whole molecule in goat. 5.0 mL Restore with deionized water (or equivalent) None 0.01% (w/v) Sodium Azide

Anti-Rabbit IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG and Rabbit Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Rabbit IgG (H&L) Secondary Antibody - Protein Information

Anti-Rabbit IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Rabbit IgG (H&L) Secondary Antibody - Images

Anti-Rabbit IgG (H&L) Secondary Antibody - Background

Anti-Rabbit IgG whole molecule antibody generated in goat detects specifically Rabbit IgG whole molecule. This secondary antibody anti-Rabbit is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.