

Human IgM (myeloma) (BULK ORDER)
Catalog # ASR3576**Specification**

Human IgM (myeloma) (BULK ORDER) - Product Information

Description	HUMAN IgM (myeloma) whole molecule (BULK ORDER)
Conjugate	Unconjugated
Physical State	Liquid (sterile filtered)
Host Isotype	IgM
Buffer	0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0
Species of Origin	Human
Preservative	0.1% (w/v) Sodium Azide

Human IgM (myeloma) (BULK ORDER) - Additional Information**Shipping Condition**

Wet Ice

Purity

Human IgM (myeloma) whole molecule was prepared from human serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Human IgM (myeloma) whole molecule was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Human Serum and anti-Human IgM (μ chain specific). No reaction was observed against anti-Human IgG F(c). Some light chain cross-reactivity will occur with anti-Human IgG.

Storage Condition

Store vial at 4° C prior to opening. Human IgM (myeloma) whole molecule is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Human IgM (myeloma) (BULK ORDER) - Protein Information**Human IgM (myeloma) (BULK ORDER) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Human IgM (myeloma) (BULK ORDER) - Images

Human IgM (myeloma) (BULK ORDER) - Background

Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum.