10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ABTB1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant ABTB1.
Catalog # AT1015a

Specification

ABTB1 Antibody (monoclonal) (M01) - Product Information

Application	**E**
Primary Accession	Q969K4
Other Accession	NM_172027
Reactivity	**Human**
Host	**mouse**
Clonality	**Monoclonal**
Isotype	**IgG2a Kappa**
Calculated MW	**53979**

ABTB1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 80325

Other Names
Ankyrin repeat and BTB/POZ domain-containing protein 1, Elongation factor 1A-binding protein, ABTB1 (HGNC:18275)

Target/Specificity
ABTB1 (NP_742024, 271 a.a. ~ 370 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format
Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage
Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions
ABTB1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

ABTB1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ABTB1 Antibody (monoclonal) (M01) - Images



Detection limit for recombinant GST tagged ABTB1 is approximately 0.03ng/ml as a capture antibody.

ABTB1 Antibody (monoclonal) (M01) - Background

This gene encodes a protein with an ankyrin repeat region and two BTB/POZ domains, which are thought to be involved in protein-protein interactions. Expression of this gene is activated by the phosphatase and tensin homolog, a tumor suppressor. Alternate splicing results in three transcript variants.

ABTB1 Antibody (monoclonal) (M01) - References

BPOZ-2 directly binds to eEF1A1 to promote eEF1A1 ubiquitylation and degradation and prevent translation. Koiwai K, et al. Genes Cells, 2008 Jun. PMID 18459963.Bood POZ containing gene type 2 is a human counterpart of yeast Btb3p and promotes the degradation of terminal deoxynucleotidyltransferase. Maezawa S, et al. Genes Cells, 2008 May. PMID 18429817.Large-scale cDNA transfection screening for genes related to cancer development and progression. Wan D, et al. Proc Natl Acad Sci U S A, 2004 Nov 2. PMID 15498874.The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.Functional proteomics mapping of a human signaling pathway. Colland F, et al. Genome Res, 2004 Jul. PMID 15231748.