

BAZ1B Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant BAZ1B. Catalog # AT1274a

Specification

BAZ1B Antibody (monoclonal) (M01) - Product Information

Application IHC, E **Primary Accession 09UIG0** Other Accession NM 023005 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 170903

BAZ1B Antibody (monoclonal) (M01) - Additional Information

Gene ID 9031

Other Names

Tyrosine-protein kinase BAZ1B, Bromodomain adjacent to zinc finger domain protein 1B, Williams syndrome transcription factor, Williams-Beuren syndrome chromosomal region 10 protein, Williams-Beuren syndrome chromosomal region 9 protein, hWALp2, BAZ1B, WBSC10, WBSCR10, WBSCR9, WSTF

Target/Specificity

BAZ1B (NP_075381, 1384 a.a. \sim 1483 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

BAZ1B Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

BAZ1B Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


BAZ1B Antibody (monoclonal) (M01) - Images



Immunoperoxidase of monoclonal antibody to BAZ1B on formalin-fixed paraffin-embedded human liver. [antibody concentration 3 ug/ml]




Detection limit for recombinant GST tagged BAZ1B is 0.3 ng/ml as a capture antibody.



Proximity Ligation Analysis of protein-protein interactions between SMARCB1 and BAZ1B HeLa cells were stained with anti-SMARCB1 rabbit purified polyclonal 1:1200 and anti-BAZ1B mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).





Proximity Ligation Analysis of protein-protein interactions between SMARCB1 and BAZ1B. Huh7 cells were stained with anti-SMARCB1 rabbit purified polyclonal 1:1200 and anti-BAZ1B mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

BAZ1B Antibody (monoclonal) (M01) - Background

This gene encodes a member of the bromodomain protein family. The bromodomain is a structural motif characteristic of proteins involved in chromatin-dependent regulation of transcription. This gene is deleted in Williams-Beuren syndrome, a developmental disorder caused by deletion of multiple genes at 7q11.23. [provided by RefSeq]