

# BIN1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant BIN1. Catalog # AT1298a

# **Specification**

# BIN1 Antibody (monoclonal) (M01) - Product Information

**Application** WB **Primary Accession** 000499 NM 004305 Other Accession Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG1 Kappa Calculated MW 64699

# BIN1 Antibody (monoclonal) (M01) - Additional Information

### Gene ID 274

### **Other Names**

Myc box-dependent-interacting protein 1, Amphiphysin II, Amphiphysin-like protein, Box-dependent myc-interacting protein 1, Bridging integrator 1, BIN1, AMPHL

## Target/Specificity

BIN1 (NP\_004296, 355 a.a.  $\sim$  454 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

# **Dilution**

WB~~1:500~1000

## **Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2.

### Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

### **Precautions**

BIN1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

### BIN1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot




- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


# BIN1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa).



BIN1 monoclonal antibody (M01), clone 1H1 Western Blot analysis of BIN1 expression in Hela S3 NE ( (Cat # AT1298a )





Western Blot analysis of BIN1 expression in transfected 293T cell line by BIN1 monoclonal antibody (M01), clone 1H1.

Lane 1: BIN1 transfected lysate(53 KDa).

Lane 2: Non-transfected lysate.

# BIN1 Antibody (monoclonal) (M01) - Background

This gene encodes several isoforms of a nucleocytoplasmic adaptor protein, one of which was initially identified as a MYC-interacting protein with features of a tumor suppressor. Isoforms that are expressed in the central nervous system may be involved in synaptic vesicle endocytosis and may interact with dynanim, synaptojanin, endophilin, and clathrin. Isoforms that are expressed in muscle and ubiquitously expressed isoforms localize to the cytoplasm and nucleus and activate a caspase-independent apoptotic process. Studies in mouse suggest that this gene plays an important role in cardiac muscle development. Alternate splicing of the gene results in ten transcript variants encoding different isoforms. Aberrant splice variants expressed in tumor cell lines have also been described.

# BIN1 Antibody (monoclonal) (M01) - References

Genetic variation and neuroimaging measures in Alzheimer disease. Biffi A, et al. Arch Neurol, 2010 Jun. PMID 20558387.Genome-wide analysis of genetic loci associated with Alzheimer disease. Seshadri S, et al. JAMA, 2010 May 12. PMID 20460622.Association of genetic variants with hemorrhagic stroke in Japanese individuals. Yoshida T, et al. Int J Mol Med, 2010 Apr. PMID 20198315.BIN1 localizes the L-type calcium channel to cardiac T-tubules. Hong TT, et al. PLoS Biol, 2010 Feb 16. PMID 20169111.Phenotype of a patient with recessive centronuclear myopathy and a novel BIN1 mutation. Claeys KG, et al. Neurology, 2010 Feb 9. PMID 20142620.