

BMP7 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a full length recombinant BMP7.

Catalog # AT1305a

Specification

BMP7 Antibody (monoclonal) (M03) - Product Information

Application	WB, E
Primary Accession	P18075
Other Accession	BC008584
Reactivity	Human, Mouse
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	49313

BMP7 Antibody (monoclonal) (M03) - Additional Information

Gene ID 655

Other Names

Bone morphogenetic protein 7, BMP-7, Osteogenic protein 1, OP-1, Eptotermin alfa, BMP7, OP1

Target/Specificity

BMP7 (AAH08584, 1 a.a. ~ 431 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

BMP7 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


BMP7 Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


BMP7 Antibody (monoclonal) (M03) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (73.15 kDa) .




BMP7 monoclonal antibody (M03), clone S52. Western Blot analysis of BMP7 expression in human ovarian cancer.




BMP7 monoclonal antibody (M03), clone S52. Western Blot analysis of BMP7 expression in A-549 (


(Cat # AT1305a)



BMP7 monoclonal antibody (M03), clone S52. Western Blot analysis of BMP7 expression in C32 (Cat # AT1305a)




BMP7 monoclonal antibody (M03), clone S52. Western Blot analysis of BMP7 expression in Raw 264.7.



Western Blot analysis of BMP7 expression in transfected 293T cell line by BMP7 monoclonal antibody (M03), clone S52.

Lane 1: BMP7 transfected lysate(49.313 KDa).

Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged BMP7 is 10 ng/ml as a capture antibody.

BMP7 Antibody (monoclonal) (M03) - Background

The bone morphogenetic proteins (BMPs) are a family of secreted signaling molecules that can induce ectopic bone growth. Many BMPs are part of the transforming growth factor-beta (TGFB) superfamily. BMPs were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. Based on its expression early in embryogenesis, the BMP encoded by this gene has a proposed role in early development and possible bone inductive activity.

BMP7 Antibody (monoclonal) (M03) - References

An approach based on a genome-wide association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014. Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Investigation of the association of BMP gene variants with nephropathy in Type 1 diabetes mellitus. McKnight AJ, et al. Diabet Med, 2010 Jun. PMID 20546278. Expression of bone morphogenetic protein 7 in lung cancer and its biological impact on lung cancer cells. Chen J, et al. Anticancer Res, 2010 Apr. PMID 20530416. L-type voltage-dependent calcium channel alpha subunit 1C is a novel candidate gene associated with secondary hyperparathyroidism: an application of haplotype-based analysis for multiple linked single nucleotide polymorphisms. Yokoyama K, et al. Nephron Clin Pract, 2010. PMID 20424473.