

**CA1 Antibody (monoclonal) (M02)****Mouse monoclonal antibody raised against a full length recombinant CA1.****Catalog # AT1354a****Specification**

---

**CA1 Antibody (monoclonal) (M02) - Product Information**

Application	WB, E
Primary Accession	<a href="#">P00915</a>
Other Accession	<a href="#">BC027890</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	28870

**CA1 Antibody (monoclonal) (M02) - Additional Information****Gene ID** 759**Other Names**

Carbonic anhydrase 1, Carbonate dehydratase I, Carbonic anhydrase B, CAB, Carbonic anhydrase I, CA-I, CA1

**Target/Specificity**

CA1 (AAH27890, 1 a.a. ~ 261 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

CA1 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


**CA1 Antibody (monoclonal) (M02) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### CA1 Antibody (monoclonal) (M02) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (54.45 kDa) .



CA1 monoclonal antibody (M02), clone M2. Western Blot analysis of CA1 expression in human lung cancer.



Detection limit for recombinant GST tagged CA1 is approximately 0.03ng/ml as a capture antibody.

### CA1 Antibody (monoclonal) (M02) - Background

Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. CA1 is closely linked to CA2 and CA3 genes on chromosome 8, and it encodes a cytosolic protein which is found at the highest level in erythrocytes. Variants of this gene have been described in some populations. Multiple alternatively spliced variants, encoding the same protein, have been identified. Transcript variants of CA1 utilizing alternative polyA\_sites have been described in literature.

#### **CA1 Antibody (monoclonal) (M02) - References**

Diabetic retinopathy is not associated with carbonic anhydrase gene polymorphisms. Abhary S, et al. Mol Vis, 2009 Jun 13. PMID 19536309. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. Decreased total carbonic anhydrase esterase activity and decreased levels of carbonic anhydrase 1 isozyme in erythrocytes of type II diabetic patients. Gambhir KK, et al. Biochem Genet, 2007 Jun. PMID 17464559. Phosph(on)ate as a zinc-binding group in metalloenzyme inhibitors: X-ray crystal structure of the antiviral drug foscarnet complexed to human carbonic anhydrase I. Temperini C, et al. Bioorg Med Chem Lett, 2007 Apr 15. PMID 17314045. Carbonic anhydrase activators: the first X-ray crystallographic study of an adduct of isoform I. Temperini C, et al. Bioorg Med Chem Lett, 2006 Oct 1. PMID 16870440.