

DCTN2 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant DCTN2.****Catalog # AT1722a****Specification**

DCTN2 Antibody (monoclonal) (M01) - Product Information

Application	IF, WB, E
Primary Accession	Q13561
Other Accession	BC000718
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 lambda
Calculated MW	44231

DCTN2 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 10540**Other Names**

Dynactin subunit 2, 50 kDa dynein-associated polypeptide, Dynactin complex 50 kDa subunit, DCTN-50, p50 dynamitin, DCTN2, DCTN50

Target/Specificity

DCTN2 (AAH00718, 302 a.a. ~ 401 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

DCTN2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


DCTN2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DCTN2 Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to DCTN2 on MCF-7 cell . [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .



DCTN2 monoclonal antibody (M01), clone 2G7 Western Blot analysis of DCTN2 expression in

MCF-7 ((Cat # AT1722a)



Detection limit for recombinant GST tagged DCTN2 is approximately 0.3ng/ml as a capture antibody.

DCTN2 Antibody (monoclonal) (M01) - Background

This gene encodes a 50-kD subunit of dynactin, a macromolecular complex consisting of 10-11 subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. It is involved in a diverse array of cellular functions, including ER-to-Golgi transport, the centripetal movement of lysosomes and endosomes, spindle formation, chromosome movement, nuclear positioning, and axonogenesis. This subunit is present in 4-5 copies per dynactin molecule. It contains three short alpha-helical coiled-coil domains that may mediate association with self or other dynactin subunits. It may interact directly with the largest subunit (p150) of dynactin and may affix p150 in place.

DCTN2 Antibody (monoclonal) (M01) - References

Molecular and functional basis for the scaffolding role of the p50/dynamitin subunit of the microtubule-associated dynactin complex. Jacquot G, et al. J Biol Chem, 2010 Jul 23. PMID 20463029. Proteomic analysis reveals Hrs ubiquitin-interacting motif-mediated ubiquitin signaling in multiple cellular processes. Pridgeon JW, et al. FEBS J, 2009 Jan. PMID 19019082. N-terminal region of ZW10 serves not only as a determinant for localization but also as a link with dynein function. Inoue M, et al. Genes Cells, 2008 Aug. PMID 18782227. Dynamitin mutagenesis reveals protein-protein interactions important for dynactin structure. Maier KC, et al. Traffic, 2008 Apr. PMID 18182012. GSK-3beta-regulated interaction of BICD with dynein is involved in microtubule anchorage at centrosome. Fumoto K, et al. EMBO J, 2006 Dec 13. PMID 17139249.