10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FUT2 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant FUT2.
Catalog # AT2123a

Specification

FUT2 Antibody (monoclonal) (M02) - Product Information

Application	**WB**
Primary Accession	Q10981
Other Accession	NM_000511
Reactivity	**Human, Mouse**
Host	**mouse**
Clonality	**Monoclonal**
Isotype	**IgG2a Kappa**
Calculated MW	**39017**

FUT2 Antibody (monoclonal) (M02) - Additional Information

Gene ID 2524

Other Names
Galactoside 2-alpha-L-fucosyltransferase 2, Alpha(1, 2)FT 2, Fucosyltransferase 2, GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase 2, SE2, Secretor blood group alpha-2-fucosyltransferase, Secretor factor, Se, FUT2, SEC2

Target/Specificity
FUT2 (NP_000502, 51 a.a. ~ 150 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution
WB~~1:500~1000

Format
Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage
Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions
FUT2 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

FUT2 Antibody (monoclonal) (M02) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FUT2 Antibody (monoclonal) (M02) - Images




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.74 KDa) .



FUT2 monoclonal antibody (M02), clone 4C12 Western Blot analysis of FUT2 expression in NIH/3T3 ((Cat # AT2123a)

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western Blot analysis of FUT2 expression in transfected 293T cell line by FUT2 monoclonal antibody (M02), clone 4C12.

Lane 1: FUT2 transfected lysate(17.5 KDa).
Lane 2: Non-transfected lysate.

FUT2 Antibody (monoclonal) (M02) - Background

The protein encoded by this gene is a Golgi stack membrane protein that is involved in the creation of a precursor of the H antigen, which is required for the final step in the soluble A and B antigen synthesis pathway. This gene is one of two encoding the galactoside 2-L-fucosyltransferase enzyme. Two transcript variants encoding the same protein have been found for this gene.

FUT2 Antibody (monoclonal) (M02) - References

Fucosyltransferase 2 (FUT2) non-secretor status is associated with Crohn's disease. McGovern DP, et al. Hum Mol Genet, 2010 Sep 1. PMID 20570966.Population based allele frequencies of disease associated polymorphisms in the Personalized Medicine Research Project. Cross DS, et al. BMC Genet, 2010 Jun 17. PMID 20565774.[Analysis of alpha-1,2-fucosyltransferase gene mutations in a Chinese family with para-Bombay phenotype] Xu XG, et al. Zhonghua Yi Xue Yi Chuan Xue Za Zhi, 2010 Jun. PMID 20533259.Polymorphisms of Lewis and Secretor genes are related to breast cancer and metastasis in axillary lymph nodes. Teresa DB, et al. Tumour Biol, 2010 Jun 1. PMID 20514537.The relationship between Lewis/Secretor genotypes and serum carbohydrate antigen 19-9 levels in a Korean population. Park HD, et al. Korean J Lab Med, 2010 Feb. PMID 20197723.