

IBRDC2 Antibody (monoclonal) (M02)**Mouse monoclonal antibody raised against a partial recombinant IBRDC2.****Catalog # AT2466a****Specification**

IBRDC2 Antibody (monoclonal) (M02) - Product Information

Application	IF, WB, IHC
Primary Accession	Q7Z419
Other Accession	NM_182757
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	33697

IBRDC2 Antibody (monoclonal) (M02) - Additional Information**Gene ID** 255488**Other Names**

E3 ubiquitin-protein ligase RNF144B, 632-, IBR domain-containing protein 2, RING finger protein 144B, p53-inducible RING finger protein, RNF144B, IBRDC2, P53RFP

Target/Specificity

IBRDC2 (NP_877434, 1 a.a. ~ 100 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

IBRDC2 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


IBRDC2 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


IBRDC2 Antibody (monoclonal) (M02) - Images



Immunofluorescence of monoclonal antibody to IBRDC2 on HeLa cell. [antibody concentration 10 ug/ml]




IBRDC2 monoclonal antibody (M02), clone 4F1 Western Blot analysis of IBRDC2 expression in IMR-32 (Cat # L008V1).



Western Blot analysis of RNF144B expression in transfected 293T cell line by IBRDC2 monoclonal antibody (M02), clone 4F1.

Lane 1: RNF144B transfected lysate(33.6 KDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to IBRDC2 on formalin-fixed paraffin-embedded human cerebellum. [antibody concentration 1.2 ug/ml]

IBRDC2 Antibody (monoclonal) (M02) - References

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. IBRDC2, an IBR-type E3 ubiquitin ligase, is a regulatory factor for Bax and apoptosis activation. Benard G, et al. EMBO J, 2010 Apr 21. PMID 20300062. A comprehensive framework of E2-RING E3 interactions of the human ubiquitin-proteasome system. van Wijk SJ, et al. Mol Syst Biol, 2009. PMID 19690564. Genome-wide association scan of quantitative traits for attention deficit hyperactivity disorder identifies novel associations and confirms candidate gene associations. Lasky-Su J, et al. Am J Med Genet B Neuropsychiatr Genet, 2008 Dec 5. PMID 18821565. The p53-inducible E3 ubiquitin ligase p53RFP induces p53-dependent apoptosis. Huang J, et al. FEBS Lett, 2006 Feb 6. PMID 16427630.