

LGALS8 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant LGALS8. Catalog # AT2702a

Specification

LGALS8 Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** 000214 BC015818 Other Accession Reactivity Human, Mouse Host mouse Clonality Monoclonal Isotype IgG2a kappa Calculated MW 35808

LGALS8 Antibody (monoclonal) (M01) - Additional Information

Gene ID 3964

Other Names

Galectin-8, Gal-8, Po66 carbohydrate-binding protein, Po66-CBP, Prostate carcinoma tumor antigen 1, PCTA-1, LGALS8

Target/Specificity

LGALS8 (AAH15818, 1 a.a. \sim 358 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

LGALS8 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

LGALS8 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot




- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


LGALS8 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (65.12 KDa).



LGALS8 monoclonal antibody (M01), clone 3E5. Western Blot analysis of LGALS8 expression in Raw 264.7.



Detection limit for recombinant GST tagged LGALS8 is approximately 0.1ng/ml as a capture antibody.

LGALS8 Antibody (monoclonal) (M01) - Background





Tel: 858.875.1900 Fax: 858.875.1999

This gene encodes a member of the galectin family. Galectins are beta-galactoside-binding animal lectins with conserved carbohydrate recognition domains. The galectins have been implicated in many essential functions including development, differentiation, cell-cell adhesion, cell-matrix interaction, growth regulation, apoptosis, and RNA splicing. This gene is widely expressed in tumoral tissues and seems to be involved in integrin-like cell interactions. Alternatively spliced transcript variants encoding different isoforms have been identified.

LGALS8 Antibody (monoclonal) (M01) - References

Galectin-8 up-regulation during hypopharyngeal and laryngeal tumor progression and comparison with galectin-1, -3 and -7. Cludts S, et al. Anticancer Res, 2009 Dec. PMID 20044599.Crystallization and preliminary X-ray diffraction analysis of a protease-resistant mutant form of human galectin-8. Yoshida H, et al. Acta Crystallogr Sect F Struct Biol Cryst Commun, 2009 May 1. PMID 19407390. Autoantibodies against galectin-8: their specificity, association with lymphopenia in systemic lupus erythematosus and detection in rheumatoid arthritis and acute inflammation. Massardo L, et al. Lupus, 2009 May. PMID 19395456. Galectin-8 interacts with podoplanin and modulates lymphatic endothelial cell functions. Cueni LN, et al. Exp Cell Res, 2009 Jun 10. PMID 19268462. Evaluation of galectin-8 expression in thyroid tumors. Savin S, et al. Med Oncol, 2009. PMID 19009371.