

**MAP3K7IP1 Antibody (monoclonal) (M03)****Mouse monoclonal antibody raised against a partial recombinant MAP3K7IP1.****Catalog # AT2780a****Specification**

---

**MAP3K7IP1 Antibody (monoclonal) (M03) - Product Information**

Application	IF, WB, E
Primary Accession	<a href="#">Q15750</a>
Other Accession	<a href="#">NM_153497</a>
Reactivity	Human, Rat
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	54644

**MAP3K7IP1 Antibody (monoclonal) (M03) - Additional Information****Gene ID** 10454**Other Names**

TGF-beta-activated kinase 1 and MAP3K7-binding protein 1, Mitogen-activated protein kinase kinase kinase 7-interacting protein 1, TGF-beta-activated kinase 1-binding protein 1, TAK1-binding protein 1, TAB1, MAP3K7IP1

**Target/Specificity**

MAP3K7IP1 (NP\_705717.1, 3 a.a. ~ 100 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

MAP3K7IP1 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


**MAP3K7IP1 Antibody (monoclonal) (M03) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### MAP3K7IP1 Antibody (monoclonal) (M03) - Images




Immunofluorescence of monoclonal antibody to MAP3K7IP1 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.52 KDa) .




MAP3K7IP1 monoclonal antibody (M03), clone 2A12 Western Blot analysis of MAP3K7IP1 expression in HeLa (Cat # AT2780a)



MAP3K7IP1 monoclonal antibody (M03), clone 2A12. Western Blot analysis of MAP3K7IP1 expression in PC-12 (Cat # AT2780a)



Detection limit for recombinant GST tagged MAP3K7IP1 is approximately 0.3ng/ml as a capture antibody.



Proximity Ligation Analysis of protein-protein interactions between HSPA1L and MAP3K7IP1 HeLa cells were stained with anti-HSPA1L rabbit purified polyclonal 1:1200 and anti-MAP3K7IP1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

#### MAP3K7IP1 Antibody (monoclonal) (M03) - Background

The protein encoded by this gene was identified as a regulator of the MAP kinase kinase kinase MAP3K7/TAK1, which is known to mediate various intracellular signaling pathways, such as those induced by TGF beta, interleukin 1, and WNT-1. This protein interacts and thus activates TAK1 kinase. It has been shown that the C-terminal portion of this protein is sufficient for binding and

activation of TAK1, while a portion of the N-terminus acts as a dominant-negative inhibitor of TGF beta, suggesting that this protein may function as a mediator between TGF beta receptors and TAK1. This protein can also interact with and activate the mitogen-activated protein kinase 14 (MAPK14/p38alpha), and thus represents an alternative activation pathway, in addition to the MAPKK pathways, which contributes to the biological responses of MAPK14 to various stimuli. Alternatively spliced transcript variants encoding distinct isoforms have been reported.

#### **MAP3K7IP1 Antibody (monoclonal) (M03) - References**

Fucosyltransferase 2 (FUT2) non-secretor status is associated with Crohn's disease. McGovern DP, et al. Hum Mol Genet, 2010 Sep 1. PMID 20570966. Autoactivation of transforming growth factor beta-activated kinase 1 is a sequential bimolecular process. Scholz R, et al. J Biol Chem, 2010 Aug 13. PMID 20538596. Association between anti-tumour necrosis factor treatment response and genetic variants within the TLR and NF{kappa}B signalling pathways. Potter C, et al. Ann Rheum Dis, 2010 Jul. PMID 20448286. The metastasis efficiency modifier ribosomal RNA processing 1 homolog B (RRP1B) is a chromatin-associated factor. Crawford NP, et al. J Biol Chem, 2009 Oct 16. PMID 19710015. Suppression of cell invasiveness by periostin via TAB1/TAK1. Isono T, et al. Int J Oncol, 2009 Aug. PMID 19578758.