

MOAP1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant MOAP1. Catalog # AT2884a

Specification

MOAP1 Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** 096BY2 Other Accession BC015044 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG1 kappa Calculated MW 39513

MOAP1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 64112

Other Names

Modulator of apoptosis 1, MAP-1, MAP1, Paraneoplastic antigen Ma4, MOAP1, PNMA4

Target/Specificity

MOAP1 (AAH15044.1, 1 a.a. \sim 351 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

MOAP1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

MOAP1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MOAP1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (64.35 KDa) .



Detection limit for recombinant GST tagged MOAP1 is approximately 1ng/ml as a capture antibody.

MOAP1 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene was identified by its interaction with apoptosis regulator BAX protein. This protein contains a Bcl-2 homology 3 (BH3)-like motif, which is required for the association with BAX. When overexpressed, this gene has been shown to mediate caspase-dependent apoptosis.

MOAP1 Antibody (monoclonal) (M01) - References

1.AICAR induces apoptosis independently of AMPK and p53 through up-regulation of the BH3-only proteins BIM and NOXA in chronic lymphocytic leukemia cells.Santidrian AF, Gonzalez-Girones DM, Iglesias-Serret D, Coll-Mulet L, Cosialls AM, de Frias M, Campas C, Gonzalez-Barca E, Alonso E, Labi V, Viollet B, Benito A, Pons G, Villunger A, Gil J.Blood. 2010 Oct 21;116(16):3023-32. Epub 2010 Jul 27.