

# NFX1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant NFX1. Catalog # AT3039a

#### Specification

# NFX1 Antibody (monoclonal) (M01) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW IF, WB, E <u>Q12986</u> <u>NM\_002504</u> Human mouse Monoclonal IgG2a Kappa 124395

# NFX1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 4799

**Other Names** Transcriptional repressor NF-X1, 632-, Nuclear transcription factor, X box-binding protein 1, NFX1, NFX2

**Target/Specificity** NFX1 (NP\_002495, 981 a.a. ~ 1080 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** NFX1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

# NFX1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>




- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

NFX1 Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to NFX1 on HeLa cell. [antibody concentration 30 ug/ml]




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.74 KDa).

250 <b>-</b> 150 <b>-</b>	
100-	
75 -	
50 -	
37 -	
25 -	£

NFX1 monoclonal antibody (M01), clone 1D12 Western Blot analysis of NFX1 expression in Hela



### S3 NE ( (Cat # AT3039a )



Detection limit for recombinant GST tagged NFX1 is approximately 0.1ng/ml as a capture antibody.

### NFX1 Antibody (monoclonal) (M01) - Background

MHC class II gene expression is controlled primarily at the transcriptional level by transcription factors that bind to the X and Y boxes, two highly conserved elements in the proximal promoter of MHC class II genes. The protein encoded by this gene is a transcriptional repressor capable of binding to the conserved X box motif of HLA-DRA and other MHC class II genes in vitro. The protein may play a role in regulating the duration of an inflammatory response by limiting the period in which class II MHC molecules are induced by IFN-gamma. Three alternative splice variants, each of which encodes a different isoform, have been identified.

# NFX1 Antibody (monoclonal) (M01) - References

NFX1-123 increases hTERT expression and telomerase activity posttranscriptionally in human papillomavirus type 16 E6 keratinocytes. Katzenellenbogen RA, et al. J Virol, 2009 Jul. PMID 19369336.NFX1 interacts with mSin3A/histone deacetylase to repress hTERT transcription in keratinocytes. Xu M, et al. Mol Cell Biol, 2008 Aug. PMID 18505829.Distinct class of putative non-conserved promoters in humans: comparative studies of alternative promoters of human and mouse genes. Tsuritani K, et al. Genome Res, 2007 Jul. PMID 17567985.NFX1-123 and poly(A) binding proteins synergistically augment activation of telomerase in human papillomavirus type 16 E6-expressing cells. Katzenellenbogen RA, et al. J Virol, 2007 Apr. PMID 17267499.Nrf1 is targeted to the endoplasmic reticulum membrane by an N-terminal transmembrane domain. Inhibition of nuclear translocation and transacting function. Wang W, et al. J Biol Chem, 2006 Jul 14. PMID 16687406.