

**PAK3 Antibody (monoclonal) (M07)****Mouse monoclonal antibody raised against a partial recombinant PAK3.****Catalog # AT3174a****Specification**

---

**PAK3 Antibody (monoclonal) (M07) - Product Information**

Application	IF, WB
Primary Accession	<a href="#">O75914</a>
Other Accession	<a href="#">NM_002578</a>
Reactivity	Human, Mouse
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	62310

**PAK3 Antibody (monoclonal) (M07) - Additional Information****Gene ID** 5063**Other Names**

Serine/threonine-protein kinase PAK 3, Beta-PAK, Oligophrenin-3, p21-activated kinase 3, PAK-3, PAK3, OPHN3

**Target/Specificity**

PAK3 (NP\_002569, 1 a.a. ~ 90 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

PAK3 Antibody (monoclonal) (M07) is for research use only and not for use in diagnostic or therapeutic procedures.


**PAK3 Antibody (monoclonal) (M07) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### PAK3 Antibody (monoclonal) (M07) - Images



Immunofluorescence of monoclonal antibody to PAK3 on NIH/3T3 cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.53 KDa) .



PAK3 monoclonal antibody (M07), clone 1H7 Western Blot analysis of PAK3 expression in NIH/3T3

( (Cat # AT3174a )

### **PAK3 Antibody (monoclonal) (M07) - Background**

PAK proteins are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signaling. PAK proteins, a family of serine/threonine p21-activating kinases, serve as targets for the small GTP binding proteins Cdc42 and RAC and have been implicated in a wide range of biological activities. The protein encoded by this gene forms an activated complex with GTP-bound RAS-like (P21), CDC2 and RAC1 proteins which then catalyzes a variety of targets. This protein may be necessary for dendritic development and for the rapid cytoskeletal reorganization in dendritic spines associated with synaptic plasticity. Defects in this gene are the cause of non-syndromic mental retardation X-linked type 30 (MRX30), also called X-linked mental retardation type 47 (MRX47). Alternatively spliced transcript variants encoding different isoforms have been identified.

### **PAK3 Antibody (monoclonal) (M07) - References**

1.A 14-3-3<sup>γ</sup> dimer-based scaffold bridges CtBP1-S/BARS to PI(4)KIII<sup>β</sup> to regulate post-Golgi carrier formation.Valente C, Turacchio G, Mariggio S, Pagliuso A, Gaibisso R, Di Tullio G, Santoro M, Formiggini F, Spano S, Piccini D, Polishchuk RS, Colanzi A, Luini A, Corda D.Nat Cell Biol. 2012 Feb 26. doi: 10.1038/ncb2445. [Epub ahead of print]