

## PCAF Antibody (monoclonal) (M04)

Mouse monoclonal antibody raised against a partial recombinant PCAF. Catalog # AT3206a

#### Specification

## PCAF Antibody (monoclonal) (M04) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW IF, WB, E <u>092831</u> <u>BC060823</u> Human mouse Monoclonal IgG2a Kappa 93013

## PCAF Antibody (monoclonal) (M04) - Additional Information

Gene ID 8850

**Other Names** Histone acetyltransferase KAT2B, Histone acetyltransferase PCAF, Histone acetylase PCAF, Lysine acetyltransferase 2B, P300/CBP-associated factor, P/CAF, KAT2B, PCAF

**Target/Specificity** PCAF (AAH60823, 353 a.a. ~ 452 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PCAF Antibody (monoclonal) (M04) is for research use only and not for use in diagnostic or therapeutic procedures.

### PCAF Antibody (monoclonal) (M04) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot




- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PCAF Antibody (monoclonal) (M04) - Images



Immunofluorescence of monoclonal antibody to PCAF on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.63 KDa).

250 - 150 - 100 - 75 -	•+
50 <b>-</b> 37 <b>-</b>	
25 <b>-</b> 20 <b>-</b>	
15- 10-	


PCAF monoclonal antibody (M04), clone 5E11. Western Blot analysis of PCAF expression in human




## uterus myoma.



PCAF monoclonal antibody (M04), clone 5E11. Western Blot analysis of PCAF expression in LNCaP ( (Cat # AT3206a )




PCAF monoclonal antibody (M04), clone 5E11 Western Blot analysis of PCAF expression in Hela S3 NE ( (Cat # AT3206a )



PCAF monoclonal antibody (M04), clone 5E11. Western Blot analysis of PCAF expression in A-431 ( (Cat # AT3206a )





Detection limit for recombinant GST tagged PCAF is approximately 0.03ng/ml as a capture antibody.

# PCAF Antibody (monoclonal) (M04) - Background

CBP and p300 are large nuclear proteins that bind to many sequence-specific factors involved in cell growth and/or differentiation, including c-jun and the adenoviral oncoprotein E1A. The protein encoded by this gene associates with p300/CBP. It has in vitro and in vivo binding activity with CBP and p300, and competes with E1A for binding sites in p300/CBP. It has histone acetyl transferase activity with core histones and nucleosome core particles, indicating that this protein plays a direct role in transcriptional regulation.

# PCAF Antibody (monoclonal) (M04) - References

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.Acetylation of lysine 564 adjacent to the C-terminal binding protein-binding motif in EVI1 is crucial for transcriptional activation of GATA2. Shimahara A, et al. J Biol Chem, 2010 May 28. PMID 20363750.CARM1 activates myogenin gene via PCAF in the early differentiation of TPA-induced rhabdomyosarcoma-derived cells. Gao X, et al. J Cell Biochem, 2010 May. PMID 20213728.Regulation of Treg functionality by acetylation-mediated Foxp3 protein stabilization. van Loosdregt J, et al. Blood, 2010 Feb 4. PMID 19996091.Single nucleotide polymorphisms in miRNA binding sites and miRNA genes as breast/ovarian cancer risk modifiers in Jewish high-risk women. Kontorovich T, et al. Int J Cancer, 2010 Aug 1. PMID 19950226.