

**PSMA7 Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a full length recombinant PSMA7.****Catalog # AT3455a****Specification**

---

**PSMA7 Antibody (monoclonal) (M01) - Product Information**

Application	IF, IP, WB, IHC, E
Primary Accession	<a href="#">O14818</a>
Other Accession	<a href="#">BC004427</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	27887

**PSMA7 Antibody (monoclonal) (M01) - Additional Information****Gene ID** 5688**Other Names**

Proteasome subunit alpha type-7, Proteasome subunit RC6-1, Proteasome subunit XAPC7, PSMA7, HSPC

**Target/Specificity**

PSMA7 (AAH04427, 1 a.a. ~ 248 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

PSMA7 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**PSMA7 Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### PSMA7 Antibody (monoclonal) (M01) - Images




Immunofluorescence of monoclonal antibody to PSMA7 on HeLa cell. [antibody concentration 1 ~ 10 ug/ml]




Immunoprecipitation of PSMA7 transfected lysate using anti-PSMA7 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with PSMA7 MaxPab rabbit polyclonal antibody.




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (53.02 KDa) .



PSMA7 monoclonal antibody (M01), clone 1A10-3G12. Western Blot analysis of PSMA7 expression in human omentum, serous carcinoma.




PSMA7 monoclonal antibody (M01), clone 1A10-3G12 Western Blot analysis of PSMA7 expression in HeLa (Cat # AT3455a)




Western Blot analysis of PSMA7 expression in transfected 293T cell line by PSMA7 monoclonal antibody (M01), clone 1A10-3G12.


Lane 1: PSMA7 transfected lysate (Predicted MW: 27.9 KDa).  
Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to PSMA7 on formalin-fixed paraffin-embedded human lung, adenosquamous cell carcinoma. [antibody concentration 5 ug/ml]



Immunoperoxidase of monoclonal antibody to PSMA7 on formalin-fixed paraffin-embedded human colon tissue. [antibody concentration 5 ug/ml]



Detection limit for recombinant GST tagged PSMA7 is approximately 1ng/ml as a capture antibody.

### PSMA7 Antibody (monoclonal) (M01) - Background

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. This particular subunit has been shown to interact specifically with the hepatitis B virus X protein, a protein critical to viral replication. In addition, this subunit is involved in regulating hepatitis virus C internal ribosome

entry site (IRES) activity, an activity essential for viral replication. This core alpha subunit is also involved in regulating the hypoxia-inducible factor-1alpha, a transcription factor important for cellular responses to oxygen tension. Multiple isoforms of this subunit arising from alternative splicing may exist but alternative transcripts for only two isoforms have been defined. A pseudogene has been identified on chromosome 9.

#### **PSMA7 Antibody (monoclonal) (M01) - References**

1.PSMA7 inhibits the tumorigenicity of A549 human lung adenocarcinoma cells.Tan JY, Huang X, Luo YL.Mol Cell Biochem. 2012 Jul;366(1-2):131-7. Epub 2012 May 15.2.Proteomic identification of putative biomarkers of radiotherapy resistance: a possible role for the 26S proteasome?Smith L, Qutob O, Watson MB, Beavis AW, Potts D, Welham KJ, Garimella V, Lind MJ, Drew PJ, Cawkwell L.Neoplasia. 2009 Nov;11(11):1194-207.