

**PSMC4 Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a full length recombinant PSMC4.****Catalog # AT3466a****Specification**

---

**PSMC4 Antibody (monoclonal) (M01) - Product Information**

Application	IF, WB, IHC, E
Primary Accession	<a href="#">P43686</a>
Other Accession	<a href="#">BC000343</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a kappa
Calculated MW	47366

**PSMC4 Antibody (monoclonal) (M01) - Additional Information****Gene ID** 5704**Other Names**

26S protease regulatory subunit 6B, 26S proteasome AAA-ATPase subunit RPT3, MB67-interacting protein, MIP224, Proteasome 26S subunit ATPase 4, Tat-binding protein 7, TBP-7, PSMC4, MIP224, TBP7

**Target/Specificity**

PSMC4 (AAH00343, 1 a.a. ~ 418 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

PSMC4 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**PSMC4 Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## PSMC4 Antibody (monoclonal) (M01) - Images




Immunofluorescence of monoclonal antibody to PSMC4 on HepG2 cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (71.72 KDa) .




PSMC4 monoclonal antibody (M01), clone 3G8 Western Blot analysis of PSMC4 expression in HepG2 (Cat # AT3466a)




Western Blot analysis of PSMC4 expression in transfected 293T cell line by PSMC4 monoclonal antibody (M01), clone 3G8.

Lane 1: PSMC4 transfected lysate(47.4 kDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to PSMC4 on formalin-fixed paraffin-embedded human colon. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged PSMC4 is approximately 0.03ng/ml as a capture antibody.

### PSMC4 Antibody (monoclonal) (M01) - Background

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure

composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit has been shown to interact with an orphan member of the nuclear hormone receptor superfamily highly expressed in liver, and with gankyrin, a liver oncoprotein. Two transcript variants encoding different isoforms have been identified.

#### **PSMC4 Antibody (monoclonal) (M01) - References**

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. Assembly pathway of the Mammalian proteasome base subcomplex is mediated by multiple specific chaperones. Kaneko T, et al. Cell, 2009 May 29. PMID 19490896. Chaperone-mediated pathway of proteasome regulatory particle assembly. Roelofs J, et al. Nature, 2009 Jun 11. PMID 19412159. A comprehensive genetic study of the proteasomal subunit S6 ATPase in German Parkinson's disease patients. Wahl C, et al. J Neural Transm, 2008 Aug. PMID 18446261. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.

#### **PSMC4 Antibody (monoclonal) (M01) - Citations**

- [Chaperone-mediated 26S Proteasome Remodeling Facilitates Free K63 Ubiquitin Chain Production and Aggresome Clearance.](#)