

QARS Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant QARS. Catalog # AT3506a

Specification

QARS Antibody (monoclonal) (M01) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW IP, WB, IHC, E <u>P47897</u> <u>NM_005051</u> Human mouse Monoclonal IgG2b Kappa 87799

QARS Antibody (monoclonal) (M01) - Additional Information

Gene ID 5859

Other Names Glutamine--tRNA ligase, Glutaminyl-tRNA synthetase, GlnRS, QARS

Target/Specificity QARS (NP_005042, 677 a.a. ~ 775 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions QARS Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

QARS Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


QARS Antibody (monoclonal) (M01) - Images

250 - 150 - 100 -	
75 -	
50 -	
37 -	
<u> </u>	
25-	

Immunoprecipitation of QARS transfected lysate using anti-QARS monoclonal antibody and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with QARS MaxPab rabbit polyclonal antibody.




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.63 KDa).






QARS monoclonal antibody (M01), clone 5F5 Western Blot analysis of QARS expression in HeLa ((Cat # AT3506a)




Western Blot analysis of QARS expression in transfected 293T cell line by QARS monoclonal antibody (M01), clone 5F5.

Lane 1: QARS transfected lysate(87.8 KDa). Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to QARS on formalin-fixed paraffin-embedded human colon. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged QARS is approximately 1ng/ml as a capture antibody. QARS Antibody (monoclonal) (M01) - Background

Aminoacyl-tRNA synthetases catalyze the aminoacylation of tRNA by their cognate amino acid. Because of their central role in linking amino acids with nucleotide triplets contained in tRNAs,



aminoacyl-tRNA synthetases are thought to be among the first proteins that appeared in evolution. In metazoans, 9 aminoacyl-tRNA synthetases specific for glutamine (gln), glutamic acid (glu), and 7 other amino acids are associated within a multienzyme complex. Although present in eukaryotes, glutaminyl-tRNA synthetase (QARS) is absent from many prokaryotes, mitochondria, and chloroplasts, in which Gln-tRNA(Gln) is formed by transamidation of the misacylated Glu-tRNA(Gln). Glutaminyl-tRNA synthetase belongs to the class-I aminoacyl-tRNA synthetase family.

QARS Antibody (monoclonal) (M01) - References

1.Proteomic identification of putative biomarkers of radiotherapy resistance: a possible role for the 26S proteasome?Smith L, Qutob O, Watson MB, Beavis AW, Potts D, Welham KJ, Garimella V, Lind MJ, Drew PJ, Cawkwell L.Neoplasia. 2009 Nov;11(11):1194-207.