

SCGB1A1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant SCGB1A1. Catalog # AT3785a

Specification

SCGB1A1 Antibody (monoclonal) (M01) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW E <u>P11684</u> <u>BC004481</u> Human mouse Monoclonal IgG2b kappa 9994

SCGB1A1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 7356

Other Names

Uteroglobin, Clara cell phospholipid-binding protein, CCPBP, Clara cells 10 kDa secretory protein, CC10, Secretoglobin family 1A member 1, Urinary protein 1, UP-1, UP1, Urine protein 1, SCGB1A1, CC10, CCSP, UGB

Target/Specificity

SCGB1A1 (AAH04481, 1 a.a. \sim 91 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions SCGB1A1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

SCGB1A1 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence



• Immunoprecipitation

- Flow Cytomety
- <u>Cell Culture</u>

SCGB1A1 Antibody (monoclonal) (M01) - Images



Detection limit for recombinant GST tagged SCGB1A1 is approximately 0.3ng/ml as a capture antibody.

SCGB1A1 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the secretoglobin family of small secreted proteins. The encoded protein has been implicated in numerous functions including anti-inflammation, inhibition of phospholipase A2 and the sequestering of hydrophobic ligands. Defects in this gene are associated with a susceptibility to asthma.

SCGB1A1 Antibody (monoclonal) (M01) - References

Association of STR polymorphisms in CMA1 and IL-4 with asthma and atopy: The SAPALDIA Cohort. Hersberger M, et al. Hum Immunol, 2010 Aug 22. PMID 20736038.Clara cell secretory protein and surfactant protein-D do not predict bronchiolitis obliterans syndrome after lung transplantation. Paantjens AW, et al. Transplantation, 2010 Aug 15. PMID 20683434.An approach based on a genome-wide association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014.Single nucleotide polymorphisms in inflammation-related genes are associated with venous thromboembolism. Beckers MM, et al. Eur J Intern Med, 2010 Aug. PMID 20603037.Independent predictive roles of eotaxin Ala23Thr, paraoxonase 2 Ser311Cys and beta-adrenergic receptor Trp64Arg polymorphisms on cardiac disease in Type 2 Diabetes--an 8-year prospective cohort analysis of 1297 patients. Wang Y, et al. Diabet Med, 2010 Apr. PMID 20536507.