

**SMAD7 Antibody (monoclonal) (M09)****Mouse monoclonal antibody raised against a partial recombinant SMAD7.****Catalog # AT3949a****Specification**

---

**SMAD7 Antibody (monoclonal) (M09) - Product Information**

|                   |                           |
|-------------------|---------------------------|
| Application       | WB, IHC, E                |
| Primary Accession | <a href="#">O15105</a>    |
| Other Accession   | <a href="#">NM_005904</a> |
| Reactivity        | Human                     |
| Host              | Mouse                     |
| Clonality         | Monoclonal                |
| Isotype           | IgG2a Kappa               |
| Calculated MW     | 46426                     |

**SMAD7 Antibody (monoclonal) (M09) - Additional Information****Gene ID** 4092**Other Names**

Mothers against decapentaplegic homolog 7, MAD homolog 7, Mothers against DPP homolog 7, Mothers against decapentaplegic homolog 8, MAD homolog 8, Mothers against DPP homolog 8, SMAD family member 7, SMAD 7, Smad7, hSMAD7, SMAD7, MADH7, MADH8

**Target/Specificity**

SMAD7 (NP\_005895, 160 a.a. ~ 260 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

SMAD7 Antibody (monoclonal) (M09) is for research use only and not for use in diagnostic or therapeutic procedures.


**SMAD7 Antibody (monoclonal) (M09) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### SMAD7 Antibody (monoclonal) (M09) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.85 kDa) .



Immunoperoxidase of monoclonal antibody to SMAD7 on formalin-fixed paraffin-embedded human placenta. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged SMAD7 is 0.3 ng/ml as a capture antibody.

### SMAD7 Antibody (monoclonal) (M09) - Background

The protein encoded by this gene is a nuclear protein that binds the E3 ubiquitin ligase SMURF2. Upon binding, this complex translocates to the cytoplasm, where it interacts with TGF-beta receptor type-1 (TGFB1), leading to the degradation of both the encoded protein and TGFB1. Expression of this gene is induced by TGFB1. Variations in this gene are a cause of susceptibility to colorectal cancer type 3 (CRC3). Several transcript variants encoding different isoforms have been found for this gene.

#### **SMAD7 Antibody (monoclonal) (M09) - References**

1. Aberrant expression in multiple components of the transforming growth factor- $\beta$ 1-induced Smad signaling pathway during 7,12-dimethylbenz[a]anthracene-induced hamster buccal-pouch squamous-cell carcinogenesis. Chen YK, Yang SH, Huang AH, Hsue SS, Lin LM. Oral Oncol. 2011 Feb 26. [Epub ahead of print]