

TCEB2 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant TCEB2.****Catalog # AT4179a****Specification**

TCEB2 Antibody (monoclonal) (M01) - Product Information

Application	IF, WB, E
Primary Accession	Q15370
Other Accession	NM_007108
Reactivity	Human, Rat
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	13133

TCEB2 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 6923**Other Names**

Transcription elongation factor B polypeptide 2, Elongin 18 kDa subunit, Elongin-B, EloB, RNA polymerase II transcription factor SIII subunit B, SIII p18, TCEB2

Target/Specificity

TCEB2 (NP_009039, 9 a.a. ~ 118 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

TCEB2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


TCEB2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TCEB2 Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to TCEB2 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .




TCEB2 monoclonal antibody (M01), clone 6F6. Western Blot analysis of TCEB2 expression in

PC-12 ((Cat # AT4179a)



TCEB2 monoclonal antibody (M01), clone 6F6 Western Blot analysis of TCEB2 expression in HeLa S3 NE ((Cat # AT4179a)



Detection limit for recombinant GST tagged TCEB2 is approximately 0.03ng/ml as a capture antibody.

TCEB2 Antibody (monoclonal) (M01) - Background

This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on chromosomes 11 and 13.

TCEB2 Antibody (monoclonal) (M01) - References

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. VHL type 2B mutations retain VBC complex form and function. Hacker KE, et al. PLoS One, 2008. PMID 19030229. Elongin B/C recruitment regulates substrate binding by CIS. Piessevaux J, et al. J Biol Chem, 2008 Aug 1. PMID 18508766. Association of SAP130/SF3b-3 with Cullin-RING ubiquitin ligase complexes and its regulation by the COP9 signalosome. Menon S, et al. BMC Biochem, 2008 Jan 3. PMID 18173839. ASB4 is a hydroxylation substrate of FIH and promotes vascular differentiation via an oxygen-dependent mechanism. Ferguson JE 3rd, et al. Mol Cell Biol, 2007 Sep. PMID 17636018.