

TNFRSF25 Antibody (monoclonal) (M06)**Mouse monoclonal antibody raised against a partial recombinant TNFRSF25.****Catalog # AT4280a****Specification**

TNFRSF25 Antibody (monoclonal) (M06) - Product Information

Application	WB, E
Primary Accession	O93038
Other Accession	NM_003790
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	45385

TNFRSF25 Antibody (monoclonal) (M06) - Additional Information**Gene ID** 8718**Other Names**

Tumor necrosis factor receptor superfamily member 25, Apo-3, Apoptosis-inducing receptor AIR, Apoptosis-mediating receptor DR3, Apoptosis-mediating receptor TRAMP, Death receptor 3, Lymphocyte-associated receptor of death, LARD, Protein WSL, Protein WSL-1, TNFRSF25, APO3, DDR3, DR3, TNFRSF12, WSL, WSL1

Target/Specificity

TNFRSF25 (NP_003781, 28 a.a. ~ 124 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

TNFRSF25 Antibody (monoclonal) (M06) is for research use only and not for use in diagnostic or therapeutic procedures.


TNFRSF25 Antibody (monoclonal) (M06) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TNFRSF25 Antibody (monoclonal) (M06) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.41 KDa) .



Detection limit for recombinant GST tagged TNFRSF25 is approximately 1ng/ml as a capture antibody.

TNFRSF25 Antibody (monoclonal) (M06) - Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed preferentially in the tissues enriched in lymphocytes, and it may play a role in regulating lymphocyte homeostasis. This receptor has been shown to stimulate NF-kappa B activity and regulate cell apoptosis. The signal transduction of this receptor is mediated by various death domain containing adaptor proteins. Knockout studies in mice suggested the role of this gene in the removal of self-reactive T cells in the thymus. Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported, most of which are potentially secreted molecules. The alternative splicing of this gene in B and T cells encounters a programmed change upon T-cell activation, which predominantly produces full-length, membrane bound isoforms, and is thought to be involved in controlling lymphocyte proliferation induced by T-cell activation.

TNFRSF25 Antibody (monoclonal) (M06) - References

Immunology: TL1A in the inflammatory network in autoimmune diseases. Bayry J. Nat Rev

Rheumatol, 2010 Feb. PMID 20125169. Association between genetic variants in VEGF, ERCC3 and occupational benzene haematotoxicity. Hosgood HD 3rd, et al. Occup Environ Med, 2009 Dec. PMID 19773279. HLA-B8, DR3: a new risk factor for graft failure after renal transplantation in patients with underlying immunoglobulin A nephropathy. Andresdottir MB, et al. Clin Transplant, 2009 Sep-Oct. PMID 19674013. Essential role of TNF receptor superfamily 25 (TNFRSF25) in the development of allergic lung inflammation. Fang L, et al. J Exp Med, 2008 May 12. PMID 18411341. Altered gene expression of caspase-10, death receptor-3 and IGFBP-3 in preeclamptic placentas. Han JY, et al. Mol Cells, 2006 Oct 31. PMID 17085968.

TNFRSF25 Antibody (monoclonal) (M06) - Citations

- [TL1A induces the expression of TGF- \$\beta\$ -inducible gene h3 \(\$\beta\$ ig-h3\) through PKC, PI3K, and ERK in THP-1 cells.](#)