

TPBG Antibody (monoclonal) (M03)**Mouse monoclonal antibody raised against a partial recombinant TPBG.****Catalog # AT4315a****Specification**

TPBG Antibody (monoclonal) (M03) - Product Information

Application	E
Primary Accession	Q13641
Other Accession	NM_006670
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	46032

TPBG Antibody (monoclonal) (M03) - Additional Information**Gene ID** 7162**Other Names**

Trophoblast glycoprotein, 5T4 oncofetal antigen, 5T4 oncofetal trophoblast glycoprotein, 5T4 oncotrophoblast glycoprotein, M6P1, Wnt-activated inhibitory factor 1, WAIF1, TPBG, 5T4

Target/Specificity

TPBG (NP_006661, 219 a.a. ~ 328 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

TPBG Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


TPBG Antibody (monoclonal) (M03) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

TPBG Antibody (monoclonal) (M03) - Images



Detection limit for recombinant GST tagged TPBG is approximately 0.1ng/ml as a capture antibody.

TPBG Antibody (monoclonal) (M03) - Background

This gene encodes a leucine-rich transmembrane glycoprotein that may be involved in cell adhesion. The encoded protein is an oncofetal antigen that is specific to trophoblast cells. In adults this protein is highly expressed in many tumor cells and is associated with poor clinical outcome in numerous cancers. Alternate splicing in the 5' UTR results in multiple transcript variants that encode the same protein.

TPBG Antibody (monoclonal) (M03) - References

An MVA-based vaccine targeting the oncofetal antigen 5T4 in patients undergoing surgical resection of colorectal cancer liver metastases. Elkord E, et al. J Immunother, 2008 Nov-Dec. PMID 18833005. Identification and functional validation of MHC class I epitopes in the tumor-associated antigen 5T4. Shingler WH, et al. Int Immunol, 2008 Aug. PMID 18567615. Placenta-derived, cellular messenger RNA expression in the maternal blood of preeclamptic women. Okazaki S, et al. Obstet Gynecol, 2007 Nov. PMID 17978129. CD8 T-cell recognition of human 5T4 oncofetal antigen. Smyth LJ, et al. Int J Cancer, 2006 Oct 1. PMID 16646078. Cell surface 5T4 antigen is transiently upregulated during early human embryonic stem cell differentiation: effect of 5T4 phenotype on neural lineage formation. Ward CM, et al. Exp Cell Res, 2006 Jun 10. PMID 16616918.