

UNG Antibody (monoclonal) (M01)
Mouse monoclonal antibody raised against a partial recombinant UNG.
Catalog # AT4473a

Specification

UNG Antibody (monoclonal) (M01) - Product Information

Application	E
Primary Accession	P13051
Other Accession	BC050634
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	34645

UNG Antibody (monoclonal) (M01) - Additional Information

Gene ID 7374

Other Names

Uracil-DNA glycosylase {ECO:0000255|HAMAP-Rule:MF_03166}, UDG
{ECO:0000255|HAMAP-Rule:MF_03166}, 32227 {ECO:0000255|HAMAP-Rule:MF_03166}, UNG
{ECO:0000255|HAMAP-Rule:MF_03166}

Target/Specificity

UNG (AAH50634, 86 a.a. ~ 190 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

UNG Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

UNG Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UNG Antibody (monoclonal) (M01) - Images

UNG Antibody (monoclonal) (M01) - Background

This gene encodes one of several uracil-DNA glycosylases. One important function of uracil-DNA glycosylases is to prevent mutagenesis by eliminating uracil from DNA molecules by cleaving the N-glycosylic bond and initiating the base-excision repair (BER) pathway. Uracil bases occur from cytosine deamination or misincorporation of dUMP residues. Alternative promoter usage and splicing of this gene leads to two different isoforms: the mitochondrial UNG1 and the nuclear UNG2.

UNG Antibody (monoclonal) (M01) - References

Polymorphisms in the base excision repair pathway and graft-versus-host disease. Arora M, et al. Leukemia, 2010 Aug. PMID 20574454. X4 and R5 HIV-1 have distinct post-entry requirements for uracil DNA glycosylase during infection of primary cells. Jones KL, et al. J Biol Chem, 2010 Jun 11. PMID 20371602. Association between genetic variants in the base excision repair pathway and outcomes after hematopoietic cell transplantations. Thyagarajan B, et al. Biol Blood Marrow Transplant, 2010 Aug. PMID 20226869. Rotational dynamics of DNA on the nucleosome surface markedly impact accessibility to a DNA repair enzyme. Hinz JM, et al. Proc Natl Acad Sci U S A, 2010 Mar 9. PMID 20176960. Customised molecular diagnosis of primary immune deficiency disorders in New Zealand: an efficient strategy for a small developed country. Ameratunga R, et al. N Z Med J, 2009 Oct 9. PMID 19859091.