

**XBP1 Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a partial recombinant XBP1.****Catalog # AT4549a****Specification**

---

**XBP1 Antibody (monoclonal) (M01) - Product Information**

Application	IF, WB
Primary Accession	<a href="#">P17861</a>
Other Accession	<a href="#">BC012841</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	28695

**XBP1 Antibody (monoclonal) (M01) - Additional Information****Gene ID** 7494**Other Names**

X-box-binding protein 1, XBP-1, Tax-responsive element-binding protein 5, XBP1, TREB5, XBP2

**Target/Specificity**

XBP1 (AAH12841, 123 a.a. ~ 225 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

XBP1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**XBP1 Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### **XBP1 Antibody (monoclonal) (M01) - Images**



Immunofluorescence of monoclonal antibody to XBP1 on HeLa cell . [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.33 KDa) .



XBP1 monoclonal antibody (M01), clone 3F5. Western Blot analysis of XBP1 expression in Hela S3 NE ( (Cat # AT4549a )

### **XPB1 Antibody (monoclonal) (M01) - Background**

This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XPB1(S), which is the functionally active transcription factor. The isoform encoded by the unspliced mRNA, XPB1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XPB1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of XPB1 has been identified and localized to chromosome 5.

### **XPB1 Antibody (monoclonal) (M01) - References**

XPB1s levels are implicated in the biology and outcome of myeloma mediating different clinical outcomes to thalidomide-based treatments. Bagratuni T, et al. Blood, 2010 Jul 15. PMID 20421453. Induction of the unfolded protein response and cell death pathway in Alzheimer's disease, but not in aged Tg2576 mice. Lee JH, et al. Exp Mol Med, 2010 May 31. PMID 20368688. Evaluation of a combinatorial cell engineering approach to overcome apoptotic effects in XPB-1(s) expressing cells. Becker E, et al. J Biotechnol, 2010 Apr 15. PMID 19958799. Direct proteasome binding and subsequent degradation of unspliced XPB-1 prevent its intracellular aggregation. Navon A, et al. FEBS Lett, 2010 Jan 4. PMID 19941857. From sugar to fat: How the transcription factor XPB1 regulates hepatic lipogenesis. Glimcher LH, et al. Ann N Y Acad Sci, 2009 Sep. PMID 19751410.