

**ZNF654 Antibody(C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5102**

**Specification**

---

**ZNF654 Antibody(C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8IZM8</a>
Other Accession	<a href="#">NP_060763.2</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=66;M=65 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**ZNF654 Antibody(C-term) - Additional Information**

**Antigen Region**  
519-545

**Other Names**  
ZNF654; Zinc finger protein 654; Melanoma-associated antigen

**Dilution**  
WB~~1:1000

**Target/Specificity**  
This ZNF654 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 519-545 amino acids from the C-terminal region of human ZNF654.

**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**  
ZNF654 Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ZNF654 Antibody(C-term) - Protein Information**

**Name** ZNF654 ([HGNC:25612](#))

**Function**

May be involved in transcriptional regulation.

#### **Cellular Location**


Nucleus.

#### **ZNF654 Antibody(C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ZNF654 Antibody(C-term) - Images**



Western blot analysis of lysates from mouse testis, mouse spleen, human liver tissue lysate (from left to right), using ZNF654 Antibody (C-term)(Cat. #AW5102). AW5102 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

#### **ZNF654 Antibody(C-term) - Background**

ZNF654 may be involved in transcriptional regulation (By similarity).