10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FBXO28 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5112

Specification

FBXO28 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	Q9NVF7
Other Accession	Q8BIG4, Q2NL16
Reactivity	**Human, Mouse**
Predicted	**Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**H=41;M=41 KDa**
Isotype	**Rabbit IgG**
Antigen Source	**Human**

FBXO28 Antibody (C-term) - Additional Information

Gene ID 23219

Antigen Region
339-368

Other Names
F-box only protein 28;KIAA0483; F-box only protein 28

Dilution
WB~~1:1000

Target/Specificity
This FBXO28 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 339-368 amino acids from the C-terminal region of human FBXO28.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
FBXO28 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FBXO28 Antibody (C-term) - Protein Information

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Name FBXO28

Synonyms CENP-30 {ECO:0000303|PubMed:20813266}, K

Function
Probably recognizes and binds to some phosphorylated proteins and promotes their ubiquitination and degradation.


Cellular Location
Chromosome, centromere, kinetochore

FBXO28 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FBXO28 Antibody (C-term) - Images



Western blot analysis of lysates from mouse brain tissue and MCF-7 cell line (from left to right), using FBXO28 Antibody (C-term)(Cat. #AW5112). AW5112 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

FBXO28 Antibody (C-term) - Background

Probably recognizes and binds to some phosphorylated proteins and promotes their ubiquitination and degradation (By similarity).

FBXO28 Antibody (C-term) - References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Gregory S.G., et al. Nature 441:315-321(2006).
Seki N., et al. DNA Res. 4:345-349(1997).
Olsen J.V., et al. Cell 127:635-648(2006).
Dephoure N., et al. Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).