

**CABLES2 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5114**

**Specification**

---

**CABLES2 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q9BTV7</a>
Other Accession	<a href="#">Q8K3M5</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=52;M=53 KDa
Isotype	Rabbit IgG
Antigen Source	Human

**CABLES2 Antibody (Center) - Additional Information**

**Gene ID** 81928

**Antigen Region**  
182-210

**Other Names**  
CABLES2;C20orf150; CDK5 and ABL1 enzyme substrate 2; CDK5 and ABL1 enzyme substrate 2;  
Interactor with CDK3 2

**Dilution**  
WB~~1:1000

**Target/Specificity**  
This CABLES2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 182-210 amino acids from the Central region of human CABLES2.

**Format**  
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**  
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**  
CABLES2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**CABLES2 Antibody (Center) - Protein Information**

**Name** CABLES2

**Synonyms** C20orf150

**Function**


Unknown. Probably involved in G1-S cell cycle transition.

**CABLES2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CABLES2 Antibody (Center) - Images**



Western blot analysis of lysates from PC-3 cell line and mouse testis tissue(from left to right), using CABLES2Antibody(Center)(Cat. #AW5114). AW5114 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

**CABLES2 Antibody (Center) - Background**

Unknown. Probably involved in G1-S cell cycle transition.

**CABLES2 Antibody (Center) - References**

Deloukas P., et al. Nature 414:865-871(2001).  
Daub H., et al. Mol. Cell 31:438-448(2008).  
Oppermann F.S., et al. Mol. Cell. Proteomics 8:1751-1764(2009).