

ADH7 Antibody (C-Term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5184

Specification

ADH7 Antibody (C-Term) - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	P40394
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=41,42;M=40;Rat=40 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

ADH7 Antibody (C-Term) - Additional Information

Gene ID 131

Antigen Region
318-346

Other Names

ADH7; Alcohol dehydrogenase class 4 mu/sigma chain; Alcohol dehydrogenase class IV mu/sigma chain; Gastric alcohol dehydrogenase; Retinol dehydrogenase

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Target/Specificity

This ADH7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 318-346 amino acids from the C-terminal region of human ADH7.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ADH7 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

ADH7 Antibody (C-Term) - Protein Information

Name ADH7 ([HGNC:256](#))

Function

Catalyzes the NAD-dependent oxidation of all-trans-retinol, alcohol, and omega-hydroxy fatty acids and their derivatives (PubMed:[15369820](http://www.uniprot.org/citations/15369820), PubMed:[16787387](http://www.uniprot.org/citations/16787387), PubMed:[9600267](http://www.uniprot.org/citations/9600267)). Oxidizes preferentially all trans-retinol, all-trans-4-hydroxyretinol, 9-cis-retinol, 2-hexenol, and long chain omega-hydroxy fatty acids such as juniperic acid (PubMed:[15369820](http://www.uniprot.org/citations/15369820), PubMed:[16787387](http://www.uniprot.org/citations/16787387), PubMed:[9600267](http://www.uniprot.org/citations/9600267)). In vitro can also catalyzes the NADH-dependent reduction of all-trans-retinal and aldehydes and their derivatives (PubMed:[15369820](http://www.uniprot.org/citations/15369820), PubMed:[16787387](http://www.uniprot.org/citations/16787387), PubMed:[9600267](http://www.uniprot.org/citations/9600267)). Reduces preferentially all trans-retinal, all-trans-4-oxoretinal and hexanal (PubMed:[15369820](http://www.uniprot.org/citations/15369820), PubMed:[16787387](http://www.uniprot.org/citations/16787387)). Catalyzes in the oxidative direction with higher efficiency (PubMed:[16787387](http://www.uniprot.org/citations/16787387), PubMed:[15369820](http://www.uniprot.org/citations/15369820)). Therefore may participate in retinoid metabolism, fatty acid omega-oxidation, and elimination of cytotoxic aldehydes produced by lipid peroxidation (PubMed:[9600267](http://www.uniprot.org/citations/9600267), PubMed:[15369820](http://www.uniprot.org/citations/15369820), PubMed:[16787387](http://www.uniprot.org/citations/16787387)).

Cellular Location

Cytoplasm.

Tissue Location


Preferentially expressed in stomach.


ADH7 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADH7 Antibody (C-Term) - Images






Confocal immunofluorescent analysis of ADH7 Antibody (C-Term) (Cat. #AW5184) with NCI-H460 cell followed by Alexa Fluor® 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).




Western blot analysis of lysates from SW480,HepG2 cell line (from left to right), using ADH7 Antibody (C-Term)(Cat. #AW5184). AW5184 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.



ADH7 Antibody (C-Term) (Cat. #AW5184) IHC analysis in formalin fixed and paraffin embedded

lung tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ADH7 Antibody (C-Term) for immunohistochemistry. Clinical relevance has not been evaluated.



ADH7 Antibody (C-Term) (Cat. #AW5184) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ADH7 Antibody (C-Term) - Background

This gene encodes class IV alcohol dehydrogenase 7 mu or sigma subunit, which is a member of the alcohol dehydrogenase family. Members of this family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. The enzyme encoded by this gene is inefficient in ethanol oxidation, but is the most active as a retinol dehydrogenase; thus it may participate in the synthesis of retinoic acid, a hormone important for cellular differentiation. The expression of this gene is much more abundant in stomach than liver, thus differing from the other known gene family members.

ADH7 Antibody (C-Term) - References

- Kedishvili, N.Y., et al. J. Biol. Chem. 270(8):3625-3630(1995)
- Cheung, B., et al. Alcohol. Clin. Exp. Res. 19(1):185-186(1995)
- Farres, J., et al. Eur. J. Biochem. 224(2):549-557(1994)
- Pares, X., et al. FEBS Lett. 303(1):69-72(1992)